


Catalytic Enantioselective Approach to the Eudesmane Sesquiterpenoids: Total Synthesis of (+)-Carissone

Samantha R. Levine, Michael R. Krout, and Brian M. Stoltz

Org. Lett., Article ASAP

Downloaded from <http://pubs.acs.org> on December 20, 2008



More About This Article

Additional resources and features associated with this article are available within the HTML version:

- Supporting Information
- Access to high resolution figures
- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article

[View the Full Text HTML](#)



ACS Publications

High quality. High impact.

Organic Letters is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Catalytic Enantioselective Approach to the Eudesmane Sesquiterpenoids: Total Synthesis of (+)-Carissone


Samantha R. Levine, Michael R. Krout, and Brian M. Stoltz*

The Arnold and Mabel Beckman Laboratories of Chemical Synthesis, Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125

stoltz@caltech.edu

Received October 19, 2008

ABSTRACT



A catalytic enantioselective approach to the eudesmane sesquiterpenoids is reported. The strategic use of a palladium-catalyzed enantioselective alkylation of vinylogous ester substrates forged the C(10) all-carbon quaternary center. This key transformation enabled a diastereoselective olefin hydrogenation to create the *syn* stereochemistry at C(7). The devised synthetic strategy allowed for the preparation of the antibacterial agent (+)-carissone and a formal synthesis of the P/Q-type calcium channel blocker (−)- α -eudesmol.

The flowering plants of the family Asteraceae (Compositae) have many historical uses, including rubber, medicines, edible oils, vegetables, and pesticides.¹ Among these flora are a large number of species that are abundant in structurally diverse sesquiterpenoids, particularly ones that contain the eudesmane skeleton (Figure 1). Over 1000 eudesmanes have been identified from these sources, with their structures diverging based on oxygenation and oxidation patterns within the carbon framework.

This ever-growing² class of important secondary metabolites possesses a wide range of biological properties, which include plant growth inhibition, insect antifeedant, antibacterial, antifungal, and antitumor activities. Representative eudesmanes include antibacterial agents (+)-carissone (**1**)³ and (+)-3-oxocostusic acid (**2**),⁴ as well as P/Q-type calcium channel blocker (−)- α -eudesmol (**3**)⁵ (Figure 1). These

(1) For a comprehensive review of the eudesmane sesquiterpenoids of the Asteraceae family, see: Wu, Q.-X.; Shi, Y.-P.; Jia, Z.-J. *Nat. Prod. Rep.* **2006**, *23*, 699–734.





Figure 1. Representative eudesmane sesquiterpenoids.

examples typify common structural motifs within this class of sesquiterpenoids, including the C(10) all-carbon quaternary stereocenter and stereogenic C(7) substituent. The structural similarities and interesting biology associated with this class of molecules has stimulated several synthetic efforts,^{6–8} most of which employ semisynthetic or chiral pool strategies. To

date, no catalytic asymmetric approach toward these eudesmanes has been developed. Herein, we report an approach⁹ that incorporates our recent method for the catalytic asymmetric formation of enantioenriched all-carbon quaternary stereocenters into a general synthetic strategy for this class of sesquiterpenoids.

In devising a strategy for accessing the eudesmanes, we simplified our target to enone **5**, which has been utilized in the preparation of structures such as **4**,^{6d} and itself embodies many features present in various family members (cf. **5** and **1**, **2**) (Scheme 1). We envisioned that the stereochemistry

Scheme 1. Retrosynthetic Analysis of the Eudesmanes



of the C(7) substituent could arise by means of the diastereoselective hydrogenation of a substituted cyclohexene (i.e., **6**), the stereochemical outcome of which would be controlled by the C(10) quaternary stereocenter. This cyclohexene could be obtained from a ring-closing methathesis of triolefin **7**, which would be derived from an appropriately substituted α -quaternary ketone (i.e., **8**). Thus, the key control element in the design of this synthetic strategy is the C(10) quaternary

(2) Fraga, B. M. *Nat. Prod. Rep.* **2007**, *24*, 1350–1381.
 (3) For the isolation and biological activity of (+)-carissone, see: (a) Mohr, K.; Schindler, O.; Reichstein, T. *Helv. Chim. Acta* **1954**, *37*, 462–471. (b) Joshi, D. V.; Boyce, S. F. *J. Org. Chem.* **1957**, *22*, 95–97. (c) Achenbach, H.; Waibel, R.; Addae-Mensah, I. *Phytochemistry* **1985**, *24*, 2325–2328. (d) Lindsay, E. A.; Berry, Y.; Jamie, J. F.; Bremner, J. B. *Phytochemistry* **2000**, *55*, 403–406.

(4) For the isolation and biological activity of (+)-3-oxocostusic acid, see: (a) Bohlmann, F.; Jakupovic, J.; Lonitz, M. *Chem. Ber.* **1977**, *110*, 301–314. (b) Khanina, M. A.; Kulyasov, A. T.; Bagryanskaya, I. Y.; Gatalov, Y. V.; Adekenov, S. M.; Raldujin, V. A. *Chem. Nat. Compd.* **1998**, *34*, 145–147. (c) Al-Dabbas, M. M.; Hashinaga, F.; Abdelgaleil, S. A. M.; Suganuma, T.; Akiyama, K.; Hayashi, H. *J. Ethnopharmacol.* **2005**, *97*, 237–240. (d) Mohamed, A. E.-H.; Ahmed, A. A.; Wollenweber, E.; Bohm, B.; Asakawa, Y. *Chem. Pharm. Bull.* **2006**, *54*, 152–155.

(5) For the isolation and biological activity of α -eudesmol, see: (a) McQuillin, F. J.; Parrack, J. D. *J. Chem. Soc.* **1956**, 2973–2978. (b) Asakura, K.; Kanemasa, T.; Minagawa, K.; Kagawa, K.; Ninomiya, M. *Brain Res.* **1999**, *823*, 169–176. (c) Toyota, M.; Yonehara, Y.; Horibe, I.; Minagawa, K.; Asakura, Y. *Phytochemistry* **1999**, *52*, 689–694. (d) Asakura, K.; Kanemasa, T.; Minagawa, K.; Kagawa, K.; Yagami, T.; Nakajima, M.; Ninomiya, M. *Brain Res.* **2000**, *873*, 94–101.

(6) For syntheses of carissone, see: (a) Pinder, A. R.; Williams, R. A. *J. Chem. Soc.* **1963**, 2773–2778. (b) Sathe, V. M.; Rao, A. S. *Indian J. Chem.* **1971**, *9*, 95–97. (c) Kutney, J. P.; Singh, A. K. *Can. J. Chem.* **1982**, *60*, 1842–1846. (d) Wang, C.-C.; Kuoh, C.-S.; Wu, T.-S. *J. Nat. Prod.* **1996**, *59*, 409–411. (e) Aoyama, Y.; Araki, Y.; Konoike, T. *Synlett* **2001**, *9*, 1452–1454.


(7) For syntheses of 3-oxocostusic acid, see: (a) Ceccherelli, P.; Curini, M.; Marcotullio, M. C.; Rosati, O. *Tetrahedron Lett.* **1990**, *31*, 3071–3074. (b) Xiong, Z.; Yang, J.; Li, Y. *Tetrahedron: Asymmetry* **1996**, *7*, 2607–2612.

stereocenter,¹⁰ and we therefore sought to develop an efficient and selective means for the preparation of this moiety.

The enantioselective alkylation of ketone enolates is an area of intense investigation in our laboratory.¹¹ This method has resulted in the preparation of wide range of carbonyl compounds with adjacent quaternary stereocenters with high levels of selectivity and excellent yields, some of which have proved valuable in synthetic endeavors.¹² The application of α -quaternary ketones such as **8** for the devised strategy would require a carbonyl transposition (i.e., **8** \rightarrow **7**), and we therefore chose to exploit the unique properties of vinylogous esters (i.e., **8** where $R^2 = OR$) pioneered by Stork and Danheiser¹³ for this purpose.

Our initial studies for the asymmetric generation of quaternary stereocenters utilizing vinylogous ester derivatives focused on enol carbonates due to preliminary investigations^{12c,14} that have demonstrated successes for similar substrates. Exposure of allyl enol carbonate **9** to typical reaction conditions consisting of a palladium(0) catalyst and ligand (*S*)-**12**¹⁵ in toluene generated vinylogous ester (+)-**13**, albeit in variable yield and selectivity (Table 1, entry 1). Unfortunately, the instability of **9** impeded further

Table 1. Asymmetric Allylation of Vinylogous Ester Derivatives^a




entry	substrate	solvent	T (°C)	product	yield ^b (%)	ee ^c (%)
1	9	PhMe	25	13	22–61	84–88
2	10	PhMe	50	13	19 ^d	79
3	10	PhMe	80	13	86	75
4	11	PhMe	50	14	86	92
5	11	PhH	50	14	61 ^e	92
6	11	THF	50	14	88	92
7	11	dioxane	50	14	90	91

^a pmdba = bis(4-methoxybenzylidene)acetone. ^b Isolated yields. ^c Enantiomeric excess determined by chiral HPLC or SFC. ^d β -Ketoester (\pm)-**10** was recovered in 69% yield. ^e β -Ketoester (\pm)-**11** was recovered in 26% yield.

studies as these results were highly dependent on the composition of this enol carbonate.¹⁶ Given the range of substrate possibilities for this transformation,^{11d} we next focused on racemic β -ketoester (\pm)-**10**. Surprisingly, this substrate proved only modestly reactive at 50 °C, producing ketone (+)-**13** in 19% yield and 79% ee (entry 2).¹⁷ Increasing the reaction temperature to 80 °C enabled complete conversion to ketone (+)-**13**, although with slightly reduced selectivity (entry 3). As the lack of reactivity seemed to be a major complication with this substrate, we considered vinylogous thioesters (i.e., (\pm)-**11**) for their reported activa-

Scheme 2. Enantioselective Synthesis of the Eudesmane Core



tion properties.¹⁴ Indeed, racemic β -ketoester (\pm)-11 did prove more reactive and produced ketone ($+$)-14 at 50 °C in good yield and 92% ee (entry 4). A screen of solvents revealed that benzene (entry 5) and ethereal solvents (entries 6 and 7) provided similar selectivities to toluene.

With optimal conditions for the preparation of 14, we sought to demonstrate the feasibility of using this ketone for the total synthesis of (+)-carisssone (1). Accordingly, racemic β -ketoester (\pm)-11 was converted to ($-$)-14 in 85% yield¹⁸ and 92% ee using ligand (R)-12, to correlate with the natural antipode of 1 (Scheme 2). Subsequent conversion of vinylogous thioester ($-$)-14 into vinylogous ester ($-$)-15 was achieved with sodium methoxide in refluxing methanol. Exposure of the resulting vinylogous ester to the substituted allylmagnesium bromide generated from 16¹⁹ provided enone ($-$)-17 in 94% yield. The success of allylmagnesium bromide additions to vinylogous ester ($-$)-15 encouraged us to investigate similar reactions of various organometallic re-

(8) For syntheses of α -eudesmol, see: (a) Humber, D. C.; Pinder, A. R.; Williams, R. A. *J. Chem. Soc.* **1967**, 2335–2340. (b) Taber, D. F.; Saleh, S. A. *Tetrahedron Lett.* **1982**, 23, 2361–2364. (c) Schwartz, M. A.; Willbrand, A. M. *J. Org. Chem.* **1985**, 50, 1359–1365. (d) Chou, T.-S.; Lee, S.-J.; Yao, N.-K. *Tetrahedron* **1989**, 45, 4113–4124. (e) Frey, B.; Hünig, S.; Koch, M.; Reissig, H.-U. *Synlett* **1991**, 854–856. (f) Frey, B.; Schnaubelt, J.; Reissig, H.-U. *Eur. J. Org. Chem.* **1999**, 1385–1393.

(9) For a review of enantioselective methodologies inspired by target-directed synthesis, see: Mohr, J. T.; Krout, M. R.; Stoltz, B. M. *Nature* **2008**, 455, 323–332.

(10) For reviews of the synthesis of quaternary stereocenters, see: (a) Cozzi, P. G.; Hilgraf, R.; Zimmermann, N. *Eur. J. Org. Chem.* **2007**, 36, 5969–5994. (b) Trost, B. M.; Jiang, C. *Synthesis* **2006**, 369–396. (c) *Quaternary Stereocenters: Challenges and Solutions for Organic Synthesis*; Christoffers, J., Baro, A., Eds.; Wiley: Weinheim, 2005. (d) Douglas, C. J.; Overman, L. E. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, 101, 5363–5367. (e) Denissova, I.; Barriault, L. *Tetrahedron* **2003**, 59, 10105–10146.

(11) (a) Behenna, D. C.; Stoltz, B. M. *J. Am. Chem. Soc.* **2004**, 126, 15044–15055. (b) Mohr, J. T.; Behenna, D. C.; Harned, A. M.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2005**, 44, 6924–6927. (c) Seto, M.; Roizen, J. L.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2008**, 47, 6873–6876. (d) For a review of the development of the enantioselective Tsuji allylation in our laboratory and others, see: Mohr, J. T.; Stoltz, B. M. *Chem.—Asian J.* **2007**, 2, 1476–1491.

(12) (a) McFadden, R. M.; Stoltz, B. M. *J. Am. Chem. Soc.* **2006**, 128, 7738–7739. (b) Behenna, D. C.; Stockdill, J. L.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2007**, 46, 4077–4080. (c) White, D. E.; Stewart, I. C.; Grubbs, R. H.; Stoltz, B. M. *J. Am. Chem. Soc.* **2008**, 130, 810–811. (d) Enquist, J. A., Jr.; Stoltz, B. M. *Nature* **2008**, 453, 1228–1231.

(13) Stork, G.; Danheiser, R. L. *J. Org. Chem.* **1973**, 38, 1775–1776.


(14) Trost, B. M.; Bream, R. N.; Xu, J. *Angew. Chem., Int. Ed.* **2006**, 45, 3109–3112.

agents with vinylogous thioester ($-$)-14; however, several conditions provided intractable mixtures with no desired products.²⁰ Nonetheless, ring-closing metathesis of enone ($-$)-17 using Grubbs' catalyst 18²¹ efficiently prepared the desired substrate (i.e., ($-$)-19) for the diastereoselective hydrogenation. Gratifyingly, heterogeneous hydrogenation utilizing Rh/Al₂O₃ catalyst²² in methanol, followed by TBS cleavage, provided alcohol ($+$)-20 in good overall yield with excellent diastereoselectivity.²³ This notable transformation generates alcohol ($+$)-20 with the C(10) and C(7) stereocenters in the desired syn configuration required for 1. Conversion of alcohol ($+$)-20 to ester ($+$)-21 was achieved by a two-step process involving Dess–Martin oxidation,²⁴ followed by chlorite oxidation²⁵ with diazomethane workup.


The availability of ester ($+$)-21 in the desired configuration enabled preparation of (+)-carisssone (1) in short order.

(15) For the development of phosphinooxazoline (PHOX) ligands, see: (a) Helmchen, G.; Pfaltz, A. *Acc. Chem. Res.* **2000**, 33, 336–345. (b) Williams, J. M. J. *Synlett* **1996**, 705–710.

(16) Enol carbonate 9 was unstable to air, forming complex reaction mixtures that substantially affected yields and selectivities for the alkylation. Aromatic carbonate *i* was identified from this mixture. See the Supporting Information for more details.



(17) The reactivity of β -ketoester (\pm)-10 contrasts significantly with that of related derivative (\pm)-*ii*, which generates *iii* in 79% yield and 86% ee under identical conditions.



(18) β -Ketoester (\pm)-11 was recovered in 9% yield.

(19) Breit, B.; Breuninger, D. *J. Am. Chem. Soc.* **2004**, 126, 10244–10245.


(20) Studies employing either allylmagnesium bromide generated from 16, MeLi, or MeMgBr nucleophiles were unsuccessful. Difficulties with these types of additions into vinylogous thioesters have been noted (see ref 14).

(21) Scholl, M.; Ding, S.; Lee, C. W.; Grubbs, R. H. *Org. Lett.* **1999**, 1, 953–956.

(22) A variety of homogeneous and heterogeneous catalysts were screened for the hydrogenation of ($-$)-19. Of those that were reactive, most promoted undesired reactivity, including enone reduction and further decomposition. Rh/Al₂O₃ proved to be a mild catalyst at 1 atm of H₂ in MeOH for this substrate.

(23) The stereochemistry of ($+$)-20 was initially verified using NOE correlations. See the Supporting Information for details.

Diastereoselective reduction of the enone carbonyl under Luche conditions,²⁶ followed by treatment of the resulting alcohol with methylmagnesium bromide^{2f} provided diol (+)-**22**²⁷ in 73% yield (Scheme 3). The preparation of this diol



intersects Aoyama's synthesis ($-$)- α -eudesmol (**3**)^{6e} and represents a formal total synthesis. Furthermore, facile allylic oxidation with manganese dioxide^{6a} gave (+)-carissoine (**1**) having spectroscopic data (^1H NMR, ^{13}C NMR, IR, HRMS, optical rotation) identical to those reported for natural **1**.

(24) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155–4156.
 (25) (a) Lindgren, B. O.; Nilsson, T. *Acta Chem. Scand.* **1973**, *27*, 888–890. (b) Kraus, G. A.; Roth, B. *J. Org. Chem.* **1980**, *45*, 4825–4830. (c) Bal, B. S.; Childers, W. E.; Pinnick, H. W. *Tetrahedron* **1981**, *37*, 2091–2096.
 (26) Luche, J. L.; Gemal, A. L. *J. Am. Chem. Soc.* **1979**, *101*, 5848–5849.

In summary, we have described the palladium-catalyzed asymmetric alkylation of vinylogous ester substrates and demonstrated the utility of such products in a general synthetic approach for the total synthesis of the eudesmane sesquiterpenoids. Fundamental to this strategy is the use of the resulting C(10) quaternary stereocenter to control the C(7) stereochemistry via a diastereoselective hydrogenation, providing a highly selective and efficient route to the antibacterial agent (+)-carissoine (**1**). Studies to understand the interplay between substrate reactivity and selectivity for the asymmetric alkylation of vinylogous ester derivatives, as well as the use of the resulting enantioenriched products in the synthesis of other bioactive natural substances,²⁸ are currently underway.

Acknowledgment. We thank Krastina V. Petrova and Justin T. Mohr (Caltech) for helpful discussions and experimental assistance. We gratefully acknowledge the NIH-NIGMS (RO1GM080269-01), Marcella R. Bonsall and the Dalton Fund (undergraduate fellowships to S.R.L.), Eli Lilly (predoctoral fellowship to M.R.K.), Amgen, Abbott Laboratories, Boehringer-Ingelheim, Materia, Merck, Bristol-Meyers Squibb, and the California Institute of Technology for financial support.

Supporting Information Available: Experimental procedures and NMR spectra for all intermediates. These materials are available free of charge via the Internet at <http://pubs.acs.org>.

OL802409H

(27) Pinder, A. R.; Williams, R. A. *J. Chem. Soc.* **1963**, 2773–2778.
 (28) For a related study, see: Petrova, K. V.; Mohr, J. T.; Stoltz, B. M. *Org. Lett.* **2009**, *11*, ASAP (ol802410t).